

EFFECTS OF SOIL FAUNA ON EARLY-STAGE LITTER DECOMPOSITION ACROSS DIVERSE TROPICAL ECOSYSTEMS IN EAST MALAYSIA

UMAR HUSSAINI TARMIZI¹, AMIRAH ALIAS²,
SITI NORASIKIN ISMAIL¹, POH SENG CHEE², YAN PENG³,
KAI YUE³, MOHAMAD AQMAL-NASER¹, AMIRRUDIN AHMAD¹,
NORHAYATI AB MANAF¹, OLGA VINDUŠKOVÁ⁴, JAN FROUZ^{1,4,5},
MAREK SEIDENGLANZ⁶, AND PETR HEDĚNEC^{1,*}

¹ Institute of Tropical Biodiversity and Sustainable Development, Universiti Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia

² Faculty of Science and Marine Environment, Universiti Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia

³ Key Laboratory for Humid Subtropical Eco-Geographical Processes of the Ministry of Education, School of Geographical Sciences, Fujian Normal University, Shangsang Road 8, Cangshan District, Fuzhou 350007, Fujian, China

⁴ Institute of Environmental Studies, Faculty of Sciences, Charles University in Prague, Benátská 2, 128 01 Prague, Czech Republic

⁵ Institute of Soil Biology and Biogeochemistry, Biology Center of CAS, Na Sádkách 702/7, 370 05 České Budějovice, Czech Republic

⁶ Agritec Plant Research, Zemědělská 2520, 787 01 Šumperk, Czech Republic

* Corresponding author: petr.hedenecek@umt.edu.my

ABSTRACT

Litterbag studies from temperate zones have shown a significant effect of soil fauna on litter decomposition. However, understanding the decomposition dynamics in tropical regions remains limited compared to temperate regions. Here we investigated the impact of soil meso- and macrofauna on litter decomposition rates at three contrasting locations in tropical area of the Eastern part of Peninsular Malaysia (tropical forest near Tasik Kenyir, permaculture in Pahang and urban soils in Universiti Malaysia Terengganu campus). We conducted litterbag experiments with different mesh sizes and soil faunal sampling to investigate the effect of soil meso- and macrofauna on litter decomposition (Fig. 1). As decomposition is fast in the tropics, we expose litterbags for three months and collect them every month. Litter mass loss increased over time, with higher decomposition rates observed in tropical forests and permaculture compared to urban soils. Tropical forest soils host significantly more diverse communities of soil fauna than the other two sites. The principal component analyses (PCA) revealed divergence in the community structure of taxonomic and functional groups among different locations, with urban soils primarily comprising *Araneae*, *Protura*, and *Diplura*, while permaculture and tropical forests mainly consisted of *Acari* and *Collembola*. Size analyses revealed that soil macrofauna enhanced decomposition rates in permaculture, while mesofauna affected decomposition in urban soils. The C:N ratio of litter in litterbags increased after three months of incubation in permaculture and tropical forest without any significant differences among mesh sizes. Random forest analyses highlighted the importance of soil moisture and texture (content of sand, silt and clay) influencing soil biota associated with decomposition processes.

Keywords: decomposition rates; litterbags; litter quality; soil macrofauna; soil mesofauna

Introduction

Soil ecosystems host an immense diversity of organisms, ranging from microorganisms like bacteria, fungi and protists to larger soil animals such as termites and millipedes (Bardgett and van der Putten 2014). For instance, one single gram of soil harbors billions of microbial cells belonging to thousands of species (Fierer 2017). While soil microbes have received considerable attention during past decades (Fierer 2017; Mahé et al. 2017; Nilsson et al. 2018), soil animals have been often overlooked until recent years (Angst et al. 2024). Many studies have unveiled remarkable diversity and abundance of soil animals (Wall et al. 2008; Lavelle et al. 2022). For example, a single square meter of soil hosts hundreds of thousands of individuals of different groups of soil invertebrates, varying with soil type and microclimatic conditions (Petersen and Luxton 1982; Wu et al. 2011; Heděnecek et al.

2022). According to body size, we can recognize microfauna (< 0.2 mm), mesofauna (> 0.2 mm), and macrofauna (> 2 mm) (Swift et al. 1979). Soil microfauna (nematodes, tardigrades, and rotifers) contribute indirectly to litter decomposition via their grazing effects on fungal and bacterial communities (Cesarz et al. 2013; Devetter et al. 2017). In contrast, macrofauna (isopods, millipedes) and mesofauna (springtails and mites) contribute directly to litter decomposition by litter fragmentation and modifying conditions favorable for bacterial and fungal decomposers (Joly et al. 2020; Mrnka et al. 2020; Coq et al. 2022).

Despite their small size, soil animals play crucial roles in ecosystem functioning, particularly in complex processes, such as litter decomposition and nutrient cycling (Filser et al. 2016; Nielsen 2019). Litter decomposition is a fundamental process in terrestrial ecosystems, influencing soil fertility, carbon sequestration, and nutrient




Fig. 1 Design of litterbag experiment.

cycling. In terrestrial ecosystems, more than 50% of the mean annual litterfall is returned to the soil via feeding activity of soil animals (Wall et al. 2008; García-Palacios et al. 2013; Heděnc et al. 2022). However, decomposition rates strongly vary between biomes, driven by multiple factors including climate, litter and substrate quality, and abundance and diversity of soil fauna (Wall et al. 2008; Heděnc et al. 2022). Soil macrofauna, such as isopods, millipedes, earthworms, and gastropods, are important drivers of leaf litter decomposition in temperate deciduous forests, while termites, millipedes, and ants are major decomposers in tropical regions (Petersen and Luxton 1982; Heděnc et al. 2022). However, the role of fauna in litter decomposition in tropical regions remains underexplored. For example, it remains unclear how land-use changes as well as differences in soil properties affect soil fauna communities and their role in litter decomposition. Land-use changes affect soil biodiversity in numerous ways (Jiangi and Purvis 2023), including pesticide use (Badani et al. 2023) and changes in soil properties (Oppong et al. 2023).

Many studies have revealed significant effects of soil fauna on litter decomposition rates (Barajas-Guzmán and Alvarez-Sánchez 2003; Wall et al. 2008; Zhang et al. 2008; García-Palacios et al. 2013; Heděnc et al. 2022). Soil macrofauna, such as isopods, millipedes, earthworms, gastropods, cockroaches and termites are responsible for litter fragmentation and colonization by microbial decomposers such as bacteria and fungi which enhance decomposition rates (Kheirallah 2004; Joly et al. 2018; Coq et al. 2022). Litterbags accessible or non-accessible for meso- and macrofauna provide a reliable tool to measure litter mass loss over time (Kampichler and Bruckner 2009; Frouz et al. 2015; Peguero et al. 2019; Peng et al. 2022b). Despite the increased number of litterbag studies (Kandeler et al. 1999; Kampichler and Bruckner 2009; Frouz et al. 2015; Peng et al. 2022b), most of these were performed in the temperate or subtropical zone while only a few studies were performed in tropical regions, especially in Peninsular Malaysia (Yamashita and Takeda 1998).

Most litterbag studies in temperate zones used relatively long incubation periods and infrequent litterbag sampling (Kandeler et al. 1999; Kampichler and Bruck-

ner 2009; Peng et al. 2022b). If the same methodology were applied in tropical regions, there would be nothing left to study, as decomposition occurs more quickly than in temperate and subtropical regions. Significant mass loss occurs during the early period of incubation in tropical regions, which has been often overlooked in long-term studies (Yamashita and Takeda 1998; Frouz et al. 2015). Therefore, studying the early decomposition stages is crucial for capturing these rapid and distinct dynamics.

To fill this knowledge gap, we established a litterbag study to (i) assess the diversity and composition of soil meso and macrofauna in soils from various contrasting habitats with different land uses with different levels of anthropogenic pressure across peninsular Malaysia; (ii) to investigate the effect of soil meso and macrofauna on litter decomposition rates during three months of incubation and (iii) to investigate influencing factors shaping litter decomposition rates at various contrasting locations in peninsular Malaysia. We hypothesize that 1) soil in tropical rain forest will host the highest diversity and abundance of soil fauna via higher resource availability and stability. In contrast, permaculture and urban soils will host lower diversity and abundances of soil animals, due to lower availability of nutrients and higher anthropogenic disturbances; 2) we hypothesize that higher litter consumption in accessible litterbags incubated in tropical forest than in permaculture and urban soils due to stable microclimatic conditions and higher diversity of soil fauna 3) we expect that soil properties will shape litter decomposition via their effect on soil biota.

Material and Methods

Study site description

Three contrasting habitats (tropical rainforest, permaculture, and urban soil at the campus of the Universiti Malaysia Terengganu) with different land uses, vegetation covers, soil properties, and composition and diversity of soil fauna were selected in the Terengganu and Pahang states in Malaysia. The tropical rainforest is located near Kenyir Lake in the Hulu Terengganu district of

Terengganu State in Malaysia (5.145357° N, 102.760592° E). The area is characterized by dominant tree vegetation with a canopy cover of 80–90%. The dominant trees are *Dipterocarpus*, *Shorea*, *Lithocarpus*, *Pometia*, *Artocarpus*, and *Maniltoa*, while dominant understory vegetation is from the family of *Zingiberaceae* and *Annonaceae*. The permaculture is situated next to the Delimah Guest House of Taman Negara in Pahang State (4.383298° N, 102.40406° E). The permaculture area is approximately 150 m². The main crops are papayas, bananas and other vegetables such as okra and cucumbers. The permaculture is managed without any inorganic fertilizers and pesticides. The permaculture is fertilized by chicken manure every three months. The post-harvest residuals are left on the site as organic biofertilizers to support natural soil decomposers. The campus of the Universiti Malaysia Terengganu (UMT) is situated in the coastal area of Kuala Nerus district of the Terengganu State (5.407046° N, 103.092333° E). The campus has approximately 26 km² and its vegetation consists of different coastal tree species, such as *Swietenia macrophylla*, *Cyrtophyllum fragrans*, and *Casuarina equisetifolia*.

Experimental design

We used three types of litterbags with different mesh sizes – those accessible for soil mesofauna (nylon net with 2 mm mesh size), macrofauna (nylon net with 2 mm mesh size and with 2 cm holes), and litterbags made from fabric with 0.1 mm mesh size as a control to test the effect of soil animals on decomposition of leaf litter. Each bag was filled with freshly fallen litter of various tree species found on the UMT campus which are also common in tropical forests across Malaysia. The leaf litter from different trees collected in the botanical garden at the UMT campus (*Agathis*, *Artocarpus*, *Cyrtophyllum*, *Dipterocarpus*, *Mangifera*, *Melaleuca*, *Milettia*, and *Shorea*) was oven-dried at 65 °C for three days and equally mixed to include all dominant tree species characterized for both natural and anthropogenic habitats in tropical areas. The experiment was conducted at three sites, each containing three subplots. In each subplot, 6 litterbags containing 5 g of oven-dried leaf litter (dried at 60 °C for 48 hours) accessible to macrofauna, 6 accessible to mesofauna, and 6 control litterbags were deployed simultaneously. The litterbags were incubated and collected in three phases: after one month, 6 macrofauna-accessible, 6 mesofauna-accessible, and 6 control litterbags were retrieved from each subplot. After two months, another set of 6 litterbags from each category were collected, and finally, after three months, the remaining 6 litterbags of each type were retrieved. In total, 162 litterbags were retrieved (Supplementary file S1). The total mass of consumed litter was measured gravimetrically in the laboratory by subtraction of dry litter weights before and after incubation. The percentage of litter consumed, decomposition constant (k value), and effect size (based on the k value) of soil mesofauna and macrofauna were measured for

each litterbag. Three litterbags of each mesh size category were randomly taken from each location to measure litter TOC, TN and C:N ratio after three months of incubation. The same number of replicates before incubation was measured as a control.

Soil fauna sampling and identification

Soil samples for mesofauna were collected using a shovel from a 10 × 10 cm square to a depth of 5 cm. A total of three compact soil blocks were taken from each location at each of the three months of incubation, resulting in a total of 27 samples. Soil mesofauna was extracted by dry extraction method using a modified Berlese-Tullgren extractor for three days (Pande and Berthet 1973). Soil mesofauna specimens were trapped in water with detergent and further stored in 70% ethanol for later identification (Peng et al. 2022a). Soil macrofauna was collected using the hand-sorting method. At each location, a soil block (25 × 25 cm and 5 cm in depth) was collected together with soil samples for mesofauna. In total, 27 monoliths were taken from all locations and sampling times. Soil monoliths were transferred to the lab and immediately processed using hand sorting of all soil fauna in size class larger than 2 mm (Swift et al. 1979; Ruiz et al. 2008). Soil animals were classified into 16 taxonomic groups at the family, order, subclass, or class level, including *Collembola*, *Acari*, *Diplura*, *Symphyla*, *Paupoda*, *Protura*, *Lumbricidae*, *Diplopoda*, *Isoptera*, *Isopoda*, *Araneae*, *Chilopoda*, *Formicidae*, *Coleoptera*, Insect others, *Blattodea* (without *Isopodera*). Furthermore, soil fauna was classified into 7 functional groups, namely Predators, Omnivores, Saprophages: Macrofauna, Saprophages: Mesofauna, Macrofauna total, Mesofauna total and Total fauna (Swift et al. 1979; Artz et al. 2010; Filser et al. 2016; Orgiazzi et al. 2016).

Soil physico-chemical properties and litter chemistry

Soil samples were collected using a shovel from a 25 × 25 cm square to a depth of 5 cm. A total of three compact soil blocks were taken from each location. Soil moisture was measured gravimetrically, before and after drying at 65 °C for 72 hours using a drying oven at the Ecological laboratory of the Faculty of Science and Marine Environment (FSSM) at UMT. The pH was determined using a Horiba LAQUA PH210, obtained from the Plant Biotechnology Lab of FSSM, at a soil-to-water ratio of 1:5. Soil texture was determined as a percentage of sand, silt, and clay after sieving through the sieve of size 2 mm, 0.02 mm and 0.002 for sand, silt, and clay respectively. Available phosphorus (P) was analyzed using the Bray-1 extraction method with the Shimadzu UV Spectrophotometer, UV-1800 model. The percentage of organic matter (OM) in soil was measured by loss of ignition method using a muffle furnace at a temperature of 375 °C for 8 hours. Total Organic Carbon (TOC), Total Nitrogen (TN), and the Carbon-to-Nitrogen (C/N) ratio were determined for soil and selected litter samples using

the Elementar UNICUBE[®] organic elemental analyzer at the Organic Chemistry Laboratory of the Faculty of Marine Engineering and Technology at UMT, with helium as a carrier gas.

Statistical analyses

We used logarithmic regression to fit the temporal trend of mass loss for each species and mesh size across three sampling locations. The decomposition rate (decomposition constant) (k value) was calculated according to the single exponential model for comparisons among different litterbags and sampling locations (Peng et al. 2022b, 2022c). Specifically, we calculated the k value (months^{-1}) using Eq. (1):

$$k = -\frac{1}{t} \ln \left(\frac{M_t}{M_0} \right) \quad (1)$$

where M_0 is the initial litter dry mass (g) and M_t is the dry mass at sampling time t (month). Alpha diversity indices (Species Richness (S), Shannon Index (H'), and Simpson's Index (D)) based on the number of taxonomic groups were calculated using the "vegan" R package (Oksanen 2015). Richness refers to the number of different species (or taxa) present in a given sample, site, or community, without considering their abundance. The principal component analyses (PCA) were employed to visualize distribution of population density of various taxonomic and functional groups of soil fauna and soil physico-chemical properties. Random forest analyses using "rfPermute" package were used to test the importance of different physico-chemical properties on litter mass loss and decomposition rate (Liu et al. 2020). The effects of litterbag mesh size, sampling locations, sampling times and their interactions on the percentage of

mass loss were assessed using a linear regression model. The Analyses of Variance (ANOVA) were used to compare the mean percentage of mass loss, decomposition rate, alpha diversity indices and densities of taxonomic and functional groups of soil fauna between three different sampling locations, followed by post hoc test (Tukey's Honest Significant Difference) for observed means. To test how access of meso- and macrofauna modulates litter decomposition rate (k) as assessed using different litterbag mesh sizes, effect sizes were calculated as a proxy. The effect sizes for mesofauna, macrofauna, and all total soil fauna were calculated as the normalized effects using the natural log response ratio (lnRR) as follows:

$$\ln RR_{mesofauna} = \ln \left(\frac{k_{2\text{ mm}}}{k_{0.2\text{ mm}}} \right) \quad (2)$$

$$\ln RR_{macrofauna} = \ln \left(\frac{k_{2\text{ cm}}}{k_{2\text{ mm}}} \right) \quad (3)$$

$$\ln RR_{total} = \ln \left(\frac{k_{2\text{ cm}}}{k_{0.2\text{ mm}}} \right) \quad (4)$$

where $k_{0.2\text{ mm}}$, $k_{2\text{ mm}}$, and $k_{2\text{ cm}}$ are the k values in the litterbags with mesh sizes of 0.1 mm, 2 mm, and 2 cm holes, respectively. The two-way ANOVA followed by Tukey's HSD test was run to test the effect of sampling location and incubation time on response ratio. The paired t-test was used to compare litter TOC, TN and C:N ratio in litterbags before and after three months of exposure in various sampling locations. Pearson correlation coefficient was used to test correlations of soil properties with litter mass loss and decomposition rates. All statistical analyses and graphical representations were conducted using the R-Studio program (R Core Team 2025) with the assistance of the "vegan" and "ggplot2" packages (Oksanen et al. 2012; Wickham 2014).

Table 1 Soil physico-chemical properties among various sampling locations.

	Sampling locations			
	Urban soil	Permaculture	Tropical forest	ANOVA
Moisture (%)	7.0±1.0c	16.9±1.9b	26.8±2.7a	***
pH	5.9±0.2b	6.1±0.2a	6.3±0.1a	***
Conductivity (uS cm ⁻¹)	36.0±8.7c	45.7±8.8b	55.4±8.9a	**
TOC (g kg ⁻¹)	13±2c	16±3b	19±5a	***
TN (g kg ⁻¹)	0.5±0.0c	0.6±0.0b	0.7±0.0a	***
C:N ratio	26±3b	27±4a	27±5a	***
Plant available P (mg kg ⁻¹)	11.8±3.0a	8.9±1.9b	5.9±0.8c	***
C:P ratio	1100±980	1800±1100	3220±3200	ns
N:P ratio	42±10c	67±11b	118±28a	***
C:N:P ratio	2.3±1b	2.8±1.0b	4.4±0a	*
OM (%)	3.8±1.0c	5.5±1.4a	6.2±1.9a	***
Sand (%)	35.0±8.8c	45.0±11.3b	55.0±13.8a	***
Silt (%)	50.0±12.5a	37.5±9.4b	25.0±6.3c	***
Clay (%)	15.0±3.8c	17.5±4.4b	20.0±5.0a	***

One-way ANOVA; *P < 0.05, **P < 0.01, ***P < 0.001. Letters indicate statistical homogenous groups.

Results

Soil properties and soil fauna community composition

Soil physico-chemical properties differed significantly among the three sampling locations (Table 1). For instance, soil moisture, conductivity, TOC, TN, N:P, and percentage of organic matter were higher in the tropical forest, intermediate in permaculture, and lowest in ur-

ban soil. The percentage of clay particles and sand was highest in the soils of the tropical forest, intermediate in permaculture, and lowest in urban soil, while the highest proportion of silt particles was found in the urban soil. The concentration of phosphorus was highest in urban soils, intermediate in permaculture, and lowest in soils of the tropical forest. The soil C:P ratio did not show any significant differences.

Table 2 Alpha diversity of soil faunal community among various locations and sampling times.

		Richness (S)±SD	Shannon index (H')±SD	Simpson index (D)±SD
Locality	Urban soil	6±1b	1.48±0.2b	0.725±0.04b
	Permaculture	6±1b	1.30±0.1b	0.610±0.04b
	Tropical forest	8±0.4a	1.75±0.1a	0.792±0.02a
	ANOVA	*	*	**
Sampling time	Month 1	6±0.9b	1.52±0.2	0.718±0.04
	Month 2	5±0b	1.39±0.1	0.709±0.02
	Month 3	8±0a	1.62±0.1	0.700±0.06
	ANOVA	**	ns	ns

Alpha diversity indices are based on the number of taxonomic groups. Two-way ANOVA; *P < 0.05, **P < 0.01. Letters indicate statistical homogenous groups.




Fig. 2 Relative abundance of functional groups of soil fauna among different locations (A) and sampling times (B). Two-way ANOVA; *P < 0.05, **P < 0.01, ***P < 0.001.

Table 3 Abundance of taxonomic and functional groups of soil fauna (individuals per m²) among various locations and sampling times.

Group	Locations				Sampling times				
	Urban soil	Permaculture	Tropical rain forest	ANOVA	Month 1	Month 2	Month 3	ANOVA	
Collembola	33±10b	17±1.8c	84±15a	***	20±6c	74±18.9a	40±8b	***	
Acari	20±6c	46±4b	74±15a	***	27±9c	70±13a	44±9b	**	
Diplura	0±0b	0±0b	20±10a	*	20±10a	0±0b	0±0b	*	
Symphyla	0±0b	3±2a	0±0b	*	3±2a	0±0b	0±0b	*	
Paupoda	0±0b	4±2a	0±0b	*	4±2a	0±0b	0±0b	*	
Protura	7±4a	0±0b	0±0b	*	0±0b	0±0b	7±4a	*	
Lumbricidae	20±10a	19±5a	0±0b	*	0±0c	30±9a	9±5b	***	
Diplopoda	4±2	0±0	3±2		3±2	0±0	4±2		
Isoptera	0±0c	4±2b	11±3a	***	4±2b	0±0c	11±3a	***	
Isopoda	0±0b	3±2a	0±0b	*	0±0b	0±0b	3±2a	*	
Araneae	29±8a	0±0c	19±3b	***	30±9a	12±3b	6±2b	***	
Chilopoda	7±2a	0±0c	3±2b	***	0±0b	3±2a	7±2a	***	
Formicidae	20±8	124±60	50±15		30±6	40±18	124±60		
Coleoptera	3±2b	0±0b	20±6a	***	3±2	7±3	13±7		
Insect others	3±2c	13±2b	20±5a	***	7.6±2b	6.4±2b	22±4a	***	
Blattodea	4±2	8±2	3±2		7±2a	0±0b	8±2a	**	
Predators	36±6a	0±0c	23±2b	***	30±9a	16±4b	13±3b	***	
Omnivores	30±8	145±63	93±12		48±4b	53±22b	168±58a	*	
Saprophages	Macrofauna	24±9	26±7		7±3b	30±9a	27±6a	*	
	Mesofauna	60.7±16b	70±8b	177±21a	***	74±19b	144±30a	91±15b	**
Total	Macrofauna	90±10	171±68	130±8		84±14	100±17	208±60	
	Mesofauna	61±16b	70±8b	177±21a	***	74±19b	144±30a	91±15b	**
Total fauna	150±15b	242±63a	307±29a	*	158±16b	243±47a	298±53a	*	

Two-way ANOVA; *P < 0.05, **P < 0.01, ***P < 0.001. Letters indicate statistical homogenous groups.




Fig. 3 Principal component analyses (PCA) of community structure of taxonomic and functional groups of soil fauna and soil physico-chemical properties among various sites and sampling times.

The alpha diversity of soil faunal groups, based on the number of taxonomic groups, differed among various locations and sampling times (Table 2). Soils in tropical forests hosted significantly higher taxonomic richness, Shannon index, and Simpson index than those in per-

maculture and urban areas. Taxonomic richness, Shannon index, and Simpson index did not differ between urban soils and permaculture. Furthermore, the taxonomic richness of soil fauna differed significantly among sampling times, showing an increasing trend over time. In

contrast, the Shannon index and Simpson index did not show any significant differences among sampling times.

The abundance of various taxonomic and functional groups differed among locations and sampling times (Table 3; Fig. 2). For example, *Acari* exhibited the highest abundance in tropical soils, intermediate abundance in permaculture, and the lowest abundance in urban soils.

Furthermore, the abundance of *Acari* varied among sampling times. The total abundance of soil fauna was higher in tropical soils and in permaculture than in urban soils. Both soil mesofauna and saprophagous mesofauna showed significantly higher densities in tropical forests compared to permaculture and urban soil. In contrast, soil macrofauna and saprophagous macrofauna did not




Fig. 4 Percentage of litter mass loss among sampling times (A) and different mesh sizes (B) and decomposition rates among different sampling sites (C). Letters indicate statistical homogenous groups.

show any significant differences among sampling sites. However, both soil macrofauna and saprophagous macrofauna exhibited differences among sampling times.

The PCA diagram revealed divergence in the community structure of taxonomic and functional groups among different locations and sampling times (Fig. 3). In urban soils, the community of soil fauna primarily comprised *Aranea*, *Protura*, and *Diplura*, and was associated with high pH, silt content, and P content. Conversely, the structure of soil fauna in permaculture and tropical forests mainly consisted of Acari and Collembola, with their community composition associated with a high C:N:P ratio, organic matter content, and soil moisture.

Effect of soil fauna on litter decomposition

Our results revealed an increase in litter mass loss in all types of litterbags over the course of increased incubation time (Fig. 4A). The litter mass loss was lowest after one month of incubation and highest after three months. The percentage of litter mass loss differed among litterbags with varying accessibility to soil meso- and macrofauna (Fig. 4B). Litterbags accessible to macrofauna

exhibited a significant difference from control litterbags but did not show any significant differences compared to those accessible to mesofauna. Litterbags accessible to mesofauna did not show any significant difference in litter mass loss compared to control litterbags. Litterbags accessible to macrofauna in tropical forest and permaculture exhibited a higher percentage of mass loss than those in urban soils. Litter decomposition rates (k values) differed significantly among litterbags incubated at different locations (Fig. 4C). Litterbags in tropical forests and permaculture exhibited higher decomposition rates than those in urban soils. The litter chemistry in litterbags incubated for three months in different locations differed significantly among sampling locations but was not affected by mesh size. The TOC and CN ratio of litter in litterbags incubated in tropical forest and permaculture increased while urban soils showed no significant changes (Figs 5A–B). In addition, TN of litter incubated in all locations decreased significantly after three months of incubation (Fig. 5C).

Effect size measurement revealed that accessibility to soil mesofauna significantly increased decomposition




Fig. 5 Total Organic Carbon (A), total Nitrogen (B) and C:N ratio (C) of litter in litterbags incubated in different sampling locations. Asterisks indicate statistically significant differences (Paired t-test; * $P < 0.05$, ** $P < 0.01$).





Fig. 6 Effect of soil mesofauna, macrofauna and total fauna on litter decomposition rates among locations (A–C) and sampling times (D–F). Letters indicate statistical homogenous groups.

A Relative importance of soil properties to decomposition rate



B Relative importance of soil properties to percentage of litter loss




Fig. 7 Relative importance of soil physico-chemical properties to decomposition rates (A) and percentage of litter mass loss (B).

Table 4 Pearson correlation coefficients of soil physico-chemical properties and litter decomposition expressed as percentage of litter mass loss and decomposition rate.

	Percentage	Decomposition rate (k, months)
Moisture	0.33*	0.34*
pH	0.11	0.14
Conductivity	0.06	0.12
TOC	0.07	0.10
TN	0.25	0.27
CN	-0.04	-0.01
P	-0.17	-0.16
CP	0.20	0.20
NP	0.27	0.27
CNP	0.16	0.16
OM	0.13	0.16
Clay	0.35*	0.37*
Silt	-0.35*	-0.36*
Sand	0.35*	0.37*

Asterisks indicate statistically significant differences, * $P < 0.05$, Bonferroni corrections were used to adjust multiple correlations.

rates in litterbags incubated in urban soils more than in permaculture without any other significant differences (Fig. 6A). Soil macrofauna increased litter decomposition rates in permaculture compared to urban soil but did not show any differences between urban soils and tropical forest or tropical forest and permaculture (Fig. 6B). Accessibility to both macrofauna and mesofauna together did not show any effect on decomposition rates (Fig. 6C). The effect of accessibility to mesofauna or macrofauna on litter decomposition did not differ among different sampling times, however, accessibility to both mesofauna and macrofauna together had a significantly larger effect on litter decomposition three months after incubation (Figs 6D–F).

Random forest analyses revealed the importance of soil moisture, sand content, clay content, and silt content, respectively, for decomposition rates (Fig. 7A). Similarly, silt content, soil moisture, clay content, and sand content, respectively, significantly contributed to litter mass loss (Fig. 7B). Pearson correlation coefficient revealed that litter decomposition rates and the percentage of mass loss increased significantly with soil moisture and with the content of clay and sand, while they decreased with silt content (Table 4).

Discussion

Soil fauna variation across locations

Our results revealed more diverse communities of soil fauna in tropical rainforest than in permaculture and urban soils. We hypothesize that variations in the diver-

sity of soil fauna are mediated by abundant and diverse resources in tropical forest, stable environmental conditions, low levels of disturbance, and complex ecological interactions. For example, the structural complexity of rainforests creates numerous microhabitats, from the mineral soils to the forest floor, which supports different soil fauna species adapted to specific niches and thus increasing overall soil biodiversity (Korboulewsky et al. 2016). In agreement with Korboulewski et al. (2016), we suggest that soil fauna is directly affected by the physical characteristics (microhabitats) and chemical composition (resource quality) of the litter specific to each tree species. In addition, soil communities are also affected by humus characteristics which are strongly linked with the litter chemistry of aboveground vegetation and often vary at different site locations (Korboulewsky et al. 2016; Schelfhout et al. 2017).

The relative abundances of various taxonomic and functional groups differed among locations and sampling times. We expect that abundances of various taxonomic and functional groups are linked to resource availability. We suggest that high nutrient content related to rapid decomposition processes in tropical habitats further supports growth and reproduction of soil organisms (Wall et al. 2008; Zhang et al. 2008; Peguero et al. 2019; Schaefer et al. 2009). Our results indicate changes in the composition of various taxonomic and functional groups of soil fauna with significantly higher densities of soil mesofauna and saprophagous mesofauna in the tropical forest compared to permaculture and urban soil. We suggest that changes in the composition of various groups of soil fauna are driven directly by aboveground vegetation via the feeding preferences of different animals. For example, a study by Gerlach et al. (2014) revealed different feeding preferences of isopods for various native and introduced plants. Moreover, a laboratory study by Heděnc et al. (2023) showed a higher consumption rate of high-quality leaf litter consumed by soil macrofauna. Similarly, a field study by Peng et al. (2022a) from the temperate zone revealed a shift in community structure of soil meso- and macrofauna in soils beneath different tree species which also differed in litter chemistry.

Our study revealed that the community structure of soil fauna in urban soils was associated with soil pH, silt content, and P content while the structure of soil fauna in permaculture and tropical forests mainly was primarily shaped by C:N:P ratio, organic matter content, and soil moisture. In addition, community structure diverged between different sampling locations. The sampling locations showed differences in soil physico-chemical properties, as well as by vegetation cover. We expect that soil physico-chemical properties are shaped by parent material and dominant vegetation (Birkhofer et al. 2012; De Schrijver et al. 2012). We expect that soil physico-chemical properties shape various groups of soil fauna indirectly via their direct effect on soil microbiota. For example, soil pH or moisture can stimulate bacterial or fungal growth

and thus can stimulate the abundance of bacterial or fungal feeders (Rousk and Bååth 2011; Heděnc et al. 2024).

Soil fauna accessibility enhances litter decomposition

Our results revealed an increase in litter mass loss for three months of incubation. We hypothesize that soil fauna would affect litter decomposition even at the early decomposition stage within three months of litterbag exposure. Increased decomposition over time can be attributed to the progressive colonization and activity of soil animals, which intensify decomposition during time. We suggest that warm and humid tropical climate supports litter decomposition in comparison with drier and colder biomes (Wall et al. 2008; Frouz et al. 2015; Heděnc et al. 2022). We also expect that stable moisture and temperature in tropical climates can accelerate the physiological activity of soil animals as well as soil microbiota, which can potentially result in high decomposition rates (Powers et al. 2009; Peguero et al. 2019).

Our study showed that litterbags accessible to macrofauna exhibited a significantly higher litter mass loss than control litterbags but did not show any significant differences compared to those accessible to mesofauna. This indicates the crucial role of macrofauna in breaking down litter, likely due to their ability to fragment larger pieces of organic matter, thereby increasing the surface area available for microbial decomposition (García-Palacios et al. 2013; Frouz 2018; Fujii et al. 2018). Our results partly corroborate with similar litterbag studies across different habitat types with various dominant vegetation (Peguero et al. 2019; Peng et al. 2022b), however, our results showed significant litter mass loss only in litterbags accessible for macrofauna. We hypothesize that body size can affect litter decomposition rate via a higher consumption rate per body weight. For instance, a study by Ardestani et al. (2019) showed that big-size animals consumed more leaves per unit of body weight than small-sized animals.

Interestingly, litterbags in tropical forests and permaculture did not differ significantly in decomposition rates, suggesting that both environments provide favorable conditions for decomposers, possibly due to higher resource availability, moisture levels, and more stable microclimatic conditions compared to urban soils. However, our results based on effect size showed that soil macrofauna showed the highest effect on litter mass loss in permaculture. This suggests that in highly biodiverse tropical forests, other factors such as microbial activity and environmental conditions might play a more dominant role in driving decomposition (Wall et al. 2008; García-Palacios et al. 2013). In contrast, in permaculture systems, where biodiversity is managed and possibly less diverse than in natural forests, macrofauna might have a more pronounced effect.

Our effect size comparison showed that mesofauna exhibited the highest effect on litter mass loss in urban soils. This could be explained by the fact that urban soils typically have lower organic matter and nutrient levels

compared to permaculture and tropical forests as shown by our data. Therefore, the presence and activity of mesofauna can have a disproportionately large impact on litter decomposition, as these organisms become critical drivers of the process (Peguero et al. 2019). Moreover, reduced biodiversity in urban soils means that mesofauna, which might otherwise be one of many decomposer groups, plays a more central role in the decomposition process in urban soils.

Our results indicated an increase of TOC and C:N ratio of litter in litterbags after three months of incubation while TN of litter in litterbags decreased significantly after three months of incubation. This suggests that soil fauna preferred the most palatable litter with a low CN ratio and left in litterbags only litter with a high C:N ratio. In addition, the passing of consumed leaf litter through the gut system also reduces the chemistry of feces which may affect the total C:N ratio of remaining litter in litterbags.

Soil properties shape fauna-mediated litter decomposition

The random forest analyses underscored the complex interplay of soil properties in shaping decomposition processes, revealing how different soil texture components, such as clay, sand, and silt, can significantly influence the rates and efficiency of litter decomposition. High clay content, for example, has been found to enhance moisture retention in soils, which is beneficial for the activity of decomposers such as microbes and soil fauna that require moist conditions to thrive (Cortez 1998; Butterly et al. 2010). Moisture is critical for enzymatic reactions involved in the breakdown of organic matter, and adequate moisture levels can facilitate the growth and activity of decomposing organisms, thereby accelerating decomposition processes (Brockett et al. 2012).

In contrast, the presence of clay in excessive amounts can also impede soil aeration, creating anaerobic conditions that are less favorable for many decomposers. In poorly aerated soils, the activity of these microorganisms is restricted, slowing down the decomposition process (Wang et al. 2021; Qian et al. 2022). Furthermore, anaerobic conditions can lead to the production of toxic substances such as methane and hydrogen sulfide, which can inhibit the activity of soil fauna and further inhibit microbial decomposition (van Agtmaal et al. 2015). Therefore, while clay content is beneficial for moisture retention, it must be balanced to avoid negative impacts on soil aeration. Furthermore, clay soils, with their fine particles, tend to be dense and compact, which can impede the movement and penetration of larger soil organisms like macrofauna (Swift et al. 1979; Frouz et al. 2006; Ruiz et al. 2008). These conditions are limiting factors for such fauna to create and maintain their own tunnels, potentially limiting their ability to forage and thrive.

Sand, on the other hand, contributes to better aeration and drainage due to its larger particle size and greater pore spaces between particles (Lavelle et al. 2020). Soils

with higher sand content tend to be well-aerated, which supports the activity of aerobic decomposers and facilitates rapid organic matter breakdown (Coleman et al. 2004; Wang et al. 2021). However, sandy soils can suffer from poor moisture retention, especially in dry conditions, which can limit the availability of water to decomposers and slow down decomposition (Coleman et al. 2004; de Vries et al. 2012; Alster et al. 2013). Thus, the positive effects of sand on soil aeration need to be balanced with sufficient moisture retention to support continuous decomposition activity. Sandy soils, with larger particles and greater pore spaces, offer less resistance to movement and are more easily penetrated by both macrofauna and mesofauna (Swift et al. 1979; Wall et al. 2008; Xin et al. 2012). These conditions facilitate the creation of tunnels and burrows, enabling soil organisms to navigate and exploit their environment effectively.

Silt particles alter moisture retention and aeration (Coleman et al. 2004). Silt improves soil structure and thus the availability of nutrients to soil organisms. However, too much silt can lead to compaction and reduce soil pore spaces, negatively impacting both aeration and water infiltration (Hartmann et al. 2014). The optimal decomposition rates are achieved when soil texture maintains a balance between these components – clay for moisture retention, sand for aeration, and silt for nutrient availability and structure (Callesen et al. 2003; Coleman et al. 2004; Velasquez et al. 2007). This balance ensures that soil conditions remain conducive to the activity of decomposers, supporting efficient organic matter breakdown and nutrient cycling in various soil environments.

Implications, limitations and future perspective

Our study revealed that soil fauna has a large impact on litter decomposition. Efficient litter decomposition by soil fauna enhances nutrient cycling, thereby improving soil fertility and crop yields (De Vries et al. 2013; Edlinger et al. 2023). The implications of efficient litter decomposition extend beyond immediate soil fertility. The presence of a robust soil fauna community can also enhance crop resilience against pests and diseases, reducing reliance on chemical pesticides (Griffiths et al. 2000; Teixeira et al. 2019; Vannier et al. 2019; Delgado-Baquerizo et al. 2020). Furthermore, our findings emphasize the importance of understanding and conserving soil fauna diversity in urban and permaculture settings, as these ecosystems are often subject to anthropogenic disturbances which can impact soil fauna communities and, consequently, litter decomposition processes (Peng et al. 2022b). By recognizing the critical role of soil fauna in litter decomposition and ecosystem functioning, efforts can be made to integrate conservation and management practices that support soil fauna biodiversity and abundance.

We suggest that our study also includes several limitations but the methods used in our study make our results comparable to other studies. Firstly, decomposition was measured solely through mass loss from the litterbags,

which does not fully capture the underlying mechanisms driving these processes. Mass loss can result from various factors, including microbial mineralization, leaching, and the fragmentation and washing out of small organic matter particles, making it difficult to distinguish the specific contributions of these processes (Kampichler and Bruckner 2009; Frouz 2018). Additionally, the role of soil fauna in litter decomposition is complex; fauna can both accelerate and slow down microbial activity through their consumption and transformation of litter (Frouz 2018; Angst et al. 2024). Furthermore, the use of litterbags with different mesh sizes may not perfectly replicate natural conditions where litter is freely accessible to all decomposers. We suggest that our study focused on specific sites and may not capture the full range of soil fauna diversity and dynamics in tropical ecosystems. Additionally, the short-term nature of the study may not fully represent long-term trends in litter decomposition dynamics. Despite the above-mentioned limitation, our study provides new insights into the initial dynamics of decomposition that are often neglected in long-term studies. We suggest that future research should employ more comprehensive methods to highlight the intricate mechanisms of litter decomposition and to fully understand the ecological roles of soil fauna. For example, combining litterbag experiments with laboratory cutting-edge methods, such as the stable isotope method, amplicon sequencing method or shotgun metagenomic method is promising to investigate specific groups of soil biota involved in specific decomposition processes (Baldrian 2017; Sultana et al. 2019; Zhang et al. 2020).

Conclusion

Our study revealed the complex interplay between soil properties, soil fauna composition, and environmental factors in shaping litter decomposition processes. For example, the significant variation observed in soil physico-chemical properties among different sampling locations highlights the diverse environmental conditions supporting various groups of soil fauna that litter decomposition. Furthermore, our results showed the impact of soil fauna on litter decomposition, with macrofauna playing a particularly significant role. Understanding the interactions between soil fauna and environmental factors, such as soil moisture and nutrient content, can provide valuable insights for ecosystem management and conservation strategies aimed at enhancing litter decomposition and nutrient cycling in terrestrial ecosystems.

Acknowledgement

This research was supported by the Ministry of Higher Education (MOHE) of Malaysia through the Fundamental Research Grant Scheme (FRGS/1/2022/STG03/

UMT/02/7). We thank Abdul Jalil Abd Rahman for providing his permaculture garden at Delima Guest House in Pahang. This research has been done with the permission of the Ministry of Natural Resources and Environmental Sustainability and Unit Perancang Ekonomi Negeri Terengganu – permit numbers 437319 and 314371.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the author(s) used ChatGPT – OpenAI version 3.2 in order to grammar check and edit English language since author(s) are not native English speakers. After using this tool/service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the content of the publication.

REFERENCES

- Alster CJ, German DP, Lu Y, Allison SD (2013) Microbial enzymatic responses to drought and to nitrogen addition in a southern California grassland. *Soil Biol Biochem* 64: 68–79. doi: 10.1016/j.soilbio.2013.03.034.
- Angst G, Potapov A, Joly F-X, Angst Š, Frouz J, Ganault P, Eisenhauer N (2024) Conceptualizing soil fauna effects on labile and stabilized soil organic matter. *Nat Commun* 15: 5005. doi: 10.1038/s41467-024-49240-x.
- Artz R, Anastasiou D, Arrouays D, Bastos A, Bendetti A, Bispo A (2010) European atlas of soil biodiversity. doi: 10.13140/RG.2.1.3178.2880.
- Badani H, Djadjou F, Haddad FZ (2023) Effects of the herbicide glyphosate [n-(phosphonomethyl) glycine] on biodiversity and organisms in the soil – A review. *Eur J Environ Sci* 13: 5–15. doi: 10.14712/23361964.2023.1.
- Baldrian P (2017) Forest microbiome: Diversity, complexity and dynamics. *FEMS Microbiol Rev* 41: 109–130.
- Barajas-Guzmán G, Alvarez-Sánchez J (2003) The relationships between litter fauna and rates of litter decomposition in a tropical rain forest. *Appl Soil Ecol* 24: 91–100. doi: 10.1016/S0929-1393(03)00069-6.
- Bardgett RD, van der Putten WH (2014) Belowground biodiversity and ecosystem functioning. *Nature* 515: 505–511. doi: 10.1038/nature13855.
- Birkhofer K, Schöning I, Alt F, Herold N, Klarner B, Maraun M, Marhan S, Oelmann Y, Wubet T, Yurkov A, Begerow D, Berner D, Buscot F, Daniel R, Diekötter T, Ehnes RB, Erdmann G, Fischer Ch, Foessel B, Groh J, Gutknecht J, Kandeler E, Lang Ch, Lohaus G, Meyer A, Nacke H, Näther A, Overmann J, Polle A, Pollierer MM, Scheu S, Schloter M, Schulze ED, Schulze W, Weinert J, Weisser W, Wolters V, Schrumpf M (2012) General relationships between abiotic soil properties and soil biota across spatial scales and different land-use types. *PLoS One* 7: e43292.
- Brockett BFT, Prescott CE, Grayston SJ (2012) Soil moisture is the major factor influencing microbial community structure and enzyme activities across seven biogeoclimatic zones in western Canada. *Soil Biol Biochem* 44: 9–20. doi: 10.1016/j.soilbio.2011.09.003.
- Butterly CR, McNeill AM, Baldock JA, Marschner P (2010) Rapid changes in carbon and phosphorus after rewetting of dry soil. *Biol Fertil Soils* 47: 41–50. doi: 10.1007/s00374-010-0500-x.
- Callesen I, Liski J, Raulund-Rasmussen K, Olsson MT, Tau-Strand L, Vesterdal L, Westman CJ (2003) Soil carbon stores in Nordic well-drained forest soils – relationships with climate and texture class. *Glob Chang Biol* 9: 358–370.
- Cesarz S, Ruess L, Jacob M, Jacob A, Schaefer M, Scheu S (2013) Tree species diversity versus tree species identity: Driving forces in structuring forest food webs as indicated by soil nematodes. *Soil Biol Biochem* 62: 36–45. doi: 10.1016/j.soilbio.2013.02.020.
- Coleman D, Crossley D, Hendrix PF (2004) *Fundamentals of Soil Ecology: Second Edition*. Academic Press, San Diego.
- Coq S, Ganault P, Le Mer G, Nahmani J, Capowiez Y, Dignac M-F, Rumpel C, Joly F-X (2022) Faeces traits as unifying predictors of detritivore effects on organic matter turnover. *Geoderma* 422: 115940. doi: 10.1016/j.geoderma.2022.115940.
- Cortez J (1998) Field decomposition of leaf litters: relationships between decomposition rates and soil moisture, soil temperature and earthworm activity. *Soil Biol Biochem* 30: 783–793. doi: 10.1016/S0038-0717(97)00163-6.
- De Schrijver A, De Frenne P, Staelens J, Verstraeten G, Muys B, Vesterdal L, Wuyts K, Van Nevel L, Schelfhout S, De Neve S, Verheyen K (2012) Tree species traits cause divergence in soil acidification during four decades of post-agricultural forest development. *Glob Chang Biol* 18: 1127–1140.
- De Vries FT, Liiri ME, Bjornlund L, Bowker MA, Christensen S, Setälä HM, Bardgett RD (2012) Land use alters the resistance and resilience of soil food webs to drought. *Nat Clim Chang* 2: 276–280.
- De Vries FT, Thébault E, Liiri M, Birkhofer K, Tsiafouli MA, Bjornlund L, Jørgensen HB, Brady MV, Christensen S, de Ruiter PC, d'Hertefeldt T, Frouz J, Hedlund K, Hemerik L, Hol WHG, Hotes S, Mortimer SR, Setälä H, Sgardelis SP, Uteseny K, van der Putten WH, Wolters V, Bardgett RD (2013) Soil food web properties explain ecosystem services across European land use systems. *Proc Natl Acad Sci USA* 110: 14296–14301.
- Delgado-Baquerizo M, Guerra CA, Cano-Díaz C, Egidi E, Wang JT, Eisenhauer N, Singh BK, Maestre FT (2020) The proportion of soil-borne pathogens increases with warming at the global scale. *Nat Clim Chang* 10: 550–554. doi: 10.1038/s41558-020-0759-3.
- Devetter M, Háněl L, Řeháková K, Doležal J (2017) Diversity and feeding strategies of soil microfauna along elevation gradients in Himalayan cold deserts. *PLoS One* 12: e0187646.
- Eddinger A, Garland G, Banerjee S, Degruno F, Garcia-Palacios P, Herzog C, Pescador DS, Romdhane S, Ryo M, Saghai A, Hallin S, Maestre FT, Philippot L, Rillig MC, van der Heijden MGA (2023) The impact of agricultural management on soil aggregation and carbon storage is regulated by climatic thresholds across a 3000 km European gradient. *Glob Chang Biol* 29: 3177–3192.
- Fierer N (2017) Embracing the unknown: disentangling the complexities of the soil microbiome. *Nat Rev Microbiol*. doi: 10.1038/nrmicro.2017.87.
- Filser J, Faber JH, Tiunov AV, Brussaard L, Frouz J, De Deyn G, Uvarov AV, Berg MP, Lavelle P, Loreau M, Wall DH, Querner P, Eijsackers H, Jiménez JJ (2016) Soil fauna: Key to new carbon models. *Soil* 2: 565–582.
- Frouz J (2018) Effects of soil macro- and mesofauna on litter decomposition and soil organic matter stabilization. *Geoderma* 332: 161–172.
- Frouz J, Elhottová D, Kuráz V, Šourková M (2006) Effects of soil macrofauna on other soil biota and soil formation in reclaimed and unreclaimed post mining sites: Results of a field microcosm experiment. *Appl Soil Ecol* 33: 308–320. doi: 10.1016/j.apsoil.2005.11.001.

- Frouz J, Roubíčková A, Heděnc P, Tajovský K (2015) Do soil fauna really hasten litter decomposition? A meta-analysis of enclosure studies. *Eur J Soil Biol* 68: 18–24.
- Fujii S, Cornelissen JHC, Berg MP, Mori AS (2018) Tree leaf and root traits mediate soil faunal contribution to litter decomposition across an elevational gradient. *Funct Ecol* 32: 840–852.
- García-Palacios P, Maestre FT, Kattge J, Wall DH (2013) Climate and litter quality differently modulate the effects of soil fauna on litter decomposition across biomes. *Ecol Lett* 16: 1045–1053. doi: 10.1111/ele.12137.
- Gerlach A, Russell DJ, Jaeschke B, Römcke J (2014) Feeding preferences of native terrestrial isopod species (Oniscoidea, Iso-poda) for native and introduced leaf litter. *Appl Soil Ecol* 83: 95–100.
- Griffiths BS, Ritz K, Bardgett RD, Cook R, Christensen S, Ekelund F, Sørensen SJ, Bååth E, Bloem J, De Ruiter PC, Dolfing J, Nicolardot B (2000) Ecosystem response of pasture soil communities to fumigation-induced microbial diversity reductions: an examination of the biodiversity-ecosystem function relationship. *Oikos* 90: 279–294. doi: 10.1034/j.1600-0706.2000.900208.x.
- Hartmann M, Niklaus PA, Zimmermann S, Schmutz S, Kremer J, Abarenkov K, Lüscher P, Widmer F, Frey B (2014) Resistance and resilience of the forest soil microbiome to logging-associated compaction. *ISME J* 8: 226–244. doi: 10.1038/ismej.2013.141.
- Heděnc P, Almahasheer H, Lin Q, Peng Y, Zheng H, Siqueira DP, Yue K, Tarmizi HU, Aqmal-Naser M, Ismail SN, Alias A, Rousk J, Vesterdal L (2024) Tree species, mycorrhizal associations, and land-use history as drivers of cohesion in soil biota communities and microbe-fauna interactions. *For Ecol Manage* 560: 121827. doi: 10.1016/j.foreco.2024.121827.
- Heděnc P, Jiménez JJ, Moradi J, Domene X, Hackenberger D, Barot S, Frossard A, Oktaba L, Filser J, Kindlmann P, Frouz J (2022) Global distribution of soil fauna functional groups and their estimated litter consumption across biomes. *Sci Rep* 12: 17362. doi: 10.1038/s41598-022-21563-z.
- Heděnc P, Zheng H, Siqueira DP, Peng Y, Schmidt IK, Frøslev TG, Kjeller R, Li H, Frouz J, Vesterdal L (2023) Litter chemistry of common European tree species drives the feeding preference and consumption rate of soil invertebrates and shapes the diversity and structure of gut and faecal microbiomes. *Soil Biol Biochem* 177: 108918. doi: 10.1016/j.soilbio.2022.108918.
- Jiang Y, Purvis A (2023) How land use affects biodiversity: An analysis of the differences in the effects recorded on different continents. *Eur J Environ Sci* 13: 15–22. doi: 10.14712/23361964.2023.2.
- Joly FX, Coq S, Coulis M, David JF, Hättenschwiler S, Mueller CW, Prater I, Subke JA (2020) Detritivore conversion of litter into faeces accelerates organic matter turnover. *Commun Biol* 3: 600. doi: 10.1038/s42003-020-01392-4.
- Joly FX, Coq S, Coulis M, Nahmani J, Hättenschwiler S (2018) Litter conversion into detritivore faeces reshuffles the quality control over C and N dynamics during decomposition. *Funct Ecol* 32: 2605–2614.
- Kampichler C, Bruckner A (2009) The role of microarthropods in terrestrial decomposition: a meta-analysis of 40 years of litterbag studies. *Biol Rev* 84: 375–389. doi: 10.1111/j.1469-185X.2009.00078.x.
- Kandeler E, Kampichler C, Joergensen RG, Mölter K (1999) Effects of mesofauna in a spruce forest on soil microbial communities and N cycling in field mesocosms. *Soil Biol Biochem* 31: 1783–1792.
- Kheirallah AM (2004) Fragmentation of leaf litter by a natural population of the millipede *Julus scandinavicus* (Latzel 1884). *Biol Fertil Soils* 10: 202–206.
- Korboulewsky N, Perez G, Chauvat M (2016) How tree diversity affects soil fauna diversity: A review. *Soil Biol Biochem* 94: 94–106. doi: 10.1016/j.soilbio.2015.11.024.
- Lavelle P, Mathieu J, Spain A, Brown G, Fragoso C, Lapied E, De Aquino A, Barois I, Barrios E, Barros ME, Bedano JC, Blanchart E, Caulfield M, Chagueza Y, Dai J, Decaëns T, Dominguez A, Dominguez Y, Feijoo A, Folgarait P, Fonte SJ, Gorosito N, Huerta E, Jimenez JJ, Kelly C, Loranger G, Marchão R, Marichal R, Praxedes C, Rodriguez L, Rousseau G, Rousseau L, Ruiz N, Sanabria C, Suarez JC, Tondoh JE, Valença AD, Vanek SJ, Vasquez J, Velasquez E, Webster E, Zhang C (2022) Soil macroinvertebrate communities: A world-wide assessment. *Glob Ecol Biogeogr* 31: 1261–1276. doi: 10.1111/geb.13492.
- Lavelle P, Spain A, Fonte S, Bedano JC, Blanchart E, Galindo V, Grimaldi M, Jimenez JJ, Velasquez E, Zangerlé A (2020) Soil aggregation, ecosystem engineers and the C cycle. *Acta Oecol* 105: 103561. doi: 10.1016/j.actao.2020.103561.
- Liu S, Wang H, Tian P, Yao X, Sun H, Wang Q, Delgado-Baquero M (2020) Decoupled diversity patterns in bacteria and fungi across continental forest ecosystems. *Soil Biol Biochem* 144: 107783.
- Mahé F, de Vargas C, Bass D, Czech L, Stamatakis A, Lara E, Singer D, Mayor J, Bunge J, Sernaker S, Siemensmeyer T, Trautmann I, Romac S, Berney C, Kozlov A, Mitchell EAD, Seppey CVW, Egge W, Lentendu G, Wirth R, Trueba G, Dunthorn M (2017) Parasites dominate hyperdiverse soil protist communities in Neotropical rainforests. *Nat Ecol Evol* 1: 91. doi: 10.1038/s41559-017-0091.
- Mrnka L, Koukol O, Hrabal R, Novák F (2020) Interactions of saprotrophic and root symbiotic fungi control the transformation of humic substances and phosphorus in Norway spruce needle litter. *Soil Biol Biochem* 149: 107919.
- Nielsen UN (2019) Functional roles of soil fauna. In: Jansson JK, Prosser JI (eds) *Soil microbiomes and their functions*. Springer, pp 137–151.
- Nilsson RH, Anslan S, Bahram M, Wurzbacher C, Baldrian P, Tedersoo L (2018) Mycobiome diversity: high-throughput sequencing and identification of fungi. *Nat Rev Microbiol* 17: 95–109. doi: 10.1038/s41579-018-0116-y.
- Oppong JC, Kešner M, Macháčková J, Kučera J, Frouz J (2023) The effect of underground drainage on peat meadows and inactivation of the drainage in an attempt to restore these meadows, which failed as it reduced the ability of soils to retain water. *Eur J Environ Sci* 13: 65–70. doi:10.14712/23361964.2023.7.
- Oksanen AJ, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB (2012) *Vegan: Community Ecology Package*. R package version 2.0-3.
- Oksanen J (2015) *Vegan: Ecological diversity*. R package version 2.3-0.
- Orgiazzi A, Bardgett R, Barrios E, Behan-Pelletier V, Briones M, Chotte JL, et al. (2016) *Global Soil Biodiversity Atlas*. European Commission, Publications Office of the European Union, Luxembourg.
- Pande YD, Berthet P (1973) Comparison of the Tullgren funnel and soil section methods for surveying oribatid populations. *Pedobiologia* 13: 182–190. <https://www.jstor.org/stable/3543884>.
- Peguero G, Sardans J, Asensio D, Fernández-Martínez M, Gargallo-Garriga A, Grau O, Llusà J, Margalef O, Márquez L, Ogaya R, Urbina I, Courtois EA, Stahl C, Langenhove LV, Verryckct LT, Richter A, Janssens IA, Peñuelas J (2019) Nutrient scarcity strengthens soil fauna control over leaf litter decomposition in tropical rainforests. *Proc R Soc B* 286: 20190848.
- Peng Y, Holmstrup M, Kappel I, De Schrijver A, Schelfhout S, Heděnc P, Zheng H, Bachega LR, Yue K, Vesterdal L (2022a)

- Litter quality, mycorrhizal association, and soil properties regulate effects of tree species on the soil fauna community. *Geoderma* 407: 115526. doi: 10.1016/j.geoderma.2021.115526.
- Peng Y, Holmstrup M, Schmidt IK, Bachega LR, Schelfhout S, Zheng H, Hedéneć P, Yue K, Vesterdal L (2022b) Tree species identity is the predominant modulator of the effects of soil fauna on leaf litter decomposition. *For Ecol Manage* 520: 120396. doi: 10.1016/j.foreco.2022.120396.
- Peng Y, Peñuelas J, Vesterdal L, Yue K, Peguero G, Fornara DA, Hedéneć P, Steffens C, Wu F (2022c) Responses of soil fauna communities to the individual and combined effects of multiple global change factors. *Ecol Lett* 25: 1376–1389. doi: 10.1111/ele.14068.
- Petersen H, Luxton M (1982) A comparative analysis of soil fauna populations and their role in decomposition processes. *Oikos* 39: 287–388.
- Powers JS, Montgomery RA, Adair EC, Brearley FQ, DeWalt SJ, Castanho CT, Deinert E, Ganzhorn JU, Gilbert ME, González-Iturbe JA, Bunyavejchewin S, Grau RH, Harms KE, Hiremath A, Iriarte-Vivar S, Manzano E, De Oliveira AA, Poorter L, Ramanamanjato JB, Salk C, Varela A, Weiblen GD, Lerda MT (2009) Decomposition in tropical forests: a pan-tropical study of the effects of litter type, litter placement and mesofaunal exclusion across a precipitation gradient. *J Ecol* 97: 801–811. doi: 10.1111/j.1365-2745.2009.01515.x.
- Qian Z, Zhuang S, Gao J, Tang L, Harindintwali JD, Wang F (2022) Aeration increases soil bacterial diversity and nutrient transformation under mulching-induced hypoxic conditions. *Sci Total Environ* 817: 153017.
- R Core Team (2025) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- Rousk J, Bååth E (2011) Growth of saprotrophic fungi and bacteria in soil. *FEMS Microbiol Ecol* 78: 17–30.
- Ruiz N, Lavelle P, Jiménez J (2008) Soil macrofauna field manual. FAO, Rome. <https://www.fao.org/4/i0211e/i0211e.pdf>.
- Schaefer M, Migge-Kleian S, Scheu S (2009) The role of soil fauna for decomposition of plant residues. In: *Functioning and Management of European Beech Ecosystems*. *Ecol Stud* 208: 207–230.
- Schelfhout S, Mertens J, Verheyen K, Vesterdal L, Baeten L, Muys B, et al. (2017) Tree species identity shapes earthworm communities. *Forests* 8: 81. doi: 10.3390/f8030081.
- Sultana N, Zhao J, Zheng Y, Cai Y, Faheem M, Peng X, Wang W, Jia Z (2019) Stable isotope probing of active methane oxidizers in rice field soils from cold regions. *Biol Fertil Soils* 55: 243–250.
- Swift MJ, Heal OW, Anderson JM (1979) *Decomposition in terrestrial ecosystems*. University of California Press, Berkeley.
- Teixeira PJP, Colaianni NR, Fitzpatrick CR, Dangl JL (2019) Beyond pathogens: microbiota interactions with the plant immune system. *Curr Opin Microbiol* 49: 7–17.
- van Agtmaal M, van Os GJ, Gera Hol WH, Hundscheid MPJ, Runia WT, Hordijk CA, de Boer W (2015) Legacy effects of anaerobic soil disinfestation on soil bacterial community composition and production of pathogen-suppressing volatiles. *Front Microbiol* 6: JUL.
- Vannier N, Agler M, Hacquard S (2019) Microbiota-mediated disease resistance in plants. *PLoS Pathog* 15: e1007740.
- Velasquez E, Lavelle P, Andrade M (2007) GISQ, a multifunctional indicator of soil quality. *Soil Biol Biochem* 39: 3066–3080.
- Wall DH, Bradford MA, St. John MG, Trofymow JA, Behan-Pelletier V, Bignell DE, Dangerfield M, Parton WJ, Rusek J, Voigt W, Wolters V, Gardel VH, Ayuke FO, Bashford R, Beljakova OI, Bohlen PJ, Brauman A, Flemming S, Henschel JR, Johnson DL, Jones TH, Kovarova M, Kranabetter JM, Kutny L, Lin KCh, Maryati M, Masse D, Pokarzhevskii A, Rahman H, Sabará MG, Salamon JA, Swift MJ, Varela A, Vasconcelos HL, White D, Zou X (2008) Global decomposition experiment shows soil animal impacts on decomposition are climate-dependent. *Glob Chang Biol* 14: 2661–2677.
- Wang Y, Tang Y, Li M, Yuan Z (2021) Aeration rate improves the compost quality of food waste and promotes the decomposition of toxic materials in leachate by changing the bacterial community. *Biores Technol* 340: 125716.
- Wickham MH (2014) Package ‘ggplot2’. <https://cran.r-project.org/package=ggplot2>.
- Wu T, Ayres E, Bardgett RD, Wall DH, Garey JR (2011) Molecular study of worldwide distribution and diversity of soil animals. *Proc Natl Acad Sci USA* 108: 17720–17725. doi: 10.1073/pnas.1103824108.
- Xin WD, Yin XQ, Song B (2012) Contribution of soil fauna to litter decomposition in Songnen sandy lands in northeastern China. *J Arid Environ* 77: 90–95. doi: 10.1016/j.jaridenv.2011.10.001.
- Yamashita T, Takeda H (1998) Decomposition and nutrient dynamics of leaf litter in litter bags of two mesh sizes set in two dipterocarp forest sites in Peninsular Malaysia. *Pedobiologia* 42: 11–21.
- Zhang D, Hui D, Luo Y, Zhou G (2008) Rates of litter decomposition in terrestrial ecosystems: global patterns and controlling factors. *J Plant Ecol* 1: 85–93.
- Zhang L, Dumont MG, Bodelier PLE, Adams JM, He D, Chu H (2020) DNA stable-isotope probing highlights the effects of temperature on functionally active methanotrophs in natural wetlands. *Soil Biol Biochem* 149: 107954.

Supplementary file S1

	Treatment	Locality	Time	Before	After	Percentage	k
A1	2 cm	Tropical_rain_forest	1	5.0	3.73	25.4	0.29
A2	2 cm	Tropical_rain_forest	1	5.0	4.41	11.8	0.13
A3	2 cm	Tropical_rain_forest	1	5.0	3.95	21.0	0.24
A4	2 cm	Tropical_rain_forest	1	5.1	4.20	17.6	0.19
A5	2 cm	Tropical_rain_forest	1	5.0	4.15	17.0	0.19
A6	2 cm	Tropical_rain_forest	1	5.1	4.18	18.0	0.20
A7	2 mm	Tropical_rain_forest	1	5.1	4.10	19.6	0.22
A8	2 mm	Tropical_rain_forest	1	5.1	3.93	22.9	0.26
A9	2 mm	Tropical_rain_forest	1	5.0	3.85	23.0	0.26
A10	2 mm	Tropical_rain_forest	1	5.0	4.06	18.8	0.21
A11	2 mm	Tropical_rain_forest	1	5.0	4.10	18.0	0.20
A12	2 mm	Tropical_rain_forest	1	5.1	4.25	16.7	0.18
A13	0.2 mm	Tropical_rain_forest	1	5.0	4.08	18.4	0.20
A14	0.2 mm	Tropical_rain_forest	1	5.0	4.23	15.4	0.17
A15	0.2 mm	Tropical_rain_forest	1	5.0	4.32	13.6	0.15
A16	0.2 mm	Tropical_rain_forest	1	5.0	4.02	19.6	0.22
A17	0.2 mm	Tropical_rain_forest	1	5.0	4.15	17.0	0.19
A18	0.2 mm	Tropical_rain_forest	1	5.0	4.12	17.6	0.19
B1	2 cm	Urban_soil	1	5.0	4.38	12.4	0.13
B2	2 cm	Urban_soil	1	5.1	4.36	14.5	0.16
B3	2 cm	Urban_soil	1	5.1	4.37	14.3	0.15
B4	2 cm	Urban_soil	1	5.1	4.45	12.7	0.14
B5	2 cm	Urban_soil	1	5.0	4.72	5.6	0.06
B6	2 cm	Urban_soil	1	5.1	4.46	12.5	0.13
B7	2 mm	Urban_soil	1	5.1	4.47	12.4	0.13
B8	2 mm	Urban_soil	1	5.0	4.44	11.2	0.12
B9	2 mm	Urban_soil	1	5.1	4.36	14.5	0.16
B10	2 mm	Urban_soil	1	5.1	4.32	15.3	0.17
B11	2 mm	Urban_soil	1	5.1	4.55	10.8	0.11
B12	2 mm	Urban_soil	1	5.1	4.36	14.5	0.16
B13	0.2 mm	Urban_soil	1	5.1	4.34	14.9	0.16
B14	0.2 mm	Urban_soil	1	5.1	4.32	15.3	0.17
B15	0.2 mm	Urban_soil	1	5.0	4.41	11.8	0.13
B16	0.2 mm	Urban_soil	1	5.1	4.37	14.3	0.15
B17	0.2 mm	Urban_soil	1	5.1	4.52	11.4	0.12
B18	0.2 mm	Urban_soil	1	5.1	4.34	14.9	0.16
C1	2 cm	Permaculture	1	5.1	4.35	14.7	0.16
C2	2 cm	Permaculture	1	5.1	4.37	14.3	0.15
C3	2 cm	Permaculture	1	5.1	4.32	15.3	0.17
C4	2 cm	Permaculture	1	5.1	4.24	16.9	0.18
C5	2 cm	Permaculture	1	5.1	3.42	32.9	0.40
C6	2 cm	Permaculture	1	5.1	4.26	16.5	0.18
C7	2 mm	Permaculture	1	5.1	4.14	18.8	0.21

	Treatment	Locality	Time	Before	After	Percentage	k
C8	2 mm	Permaculture	1	5.1	4.19	17.8	0.20
C9	2 mm	Permaculture	1	5.1	4.27	16.3	0.18
C10	2 mm	Permaculture	1	5.0	4.30	14.0	0.15
C11	2 mm	Permaculture	1	5.0	4.28	14.4	0.16
C12	2 mm	Permaculture	1	5.0	4.29	14.2	0.15
C13	0.2 mm	Permaculture	1	5.0	4.00	20.0	0.22
C14	0.2 mm	Permaculture	1	5.1	4.38	14.1	0.15
C15	0.2 mm	Permaculture	1	5.0	4.25	15.0	0.16
C16	0.2 mm	Permaculture	1	5.0	4.08	18.4	0.20
C17	0.2 mm	Permaculture	1	5.1	4.14	18.8	0.21
C18	0.2 mm	Permaculture	1	5.1	4.4	13.7	0.15
D1	2 cm	Urban_soil	2	5.0	4.03	19.4	0.11
D2	2 cm	Urban_soil	2	5.1	4.03	21.0	0.12
D3	2 cm	Urban_soil	2	5.1	4.09	19.8	0.11
D4	2 cm	Urban_soil	2	5.0	4.09	18.2	0.10
D5	2 cm	Urban_soil	2	5.1	4.16	18.4	0.10
D6	2 cm	Urban_soil	2	5.1	4.22	17.3	0.09
D7	2 mm	Urban_soil	2	5.1	4.06	20.4	0.11
D8	2 mm	Urban_soil	2	5.1	4.16	18.4	0.10
D9	2 mm	Urban_soil	2	5.1	4.05	20.6	0.12
D10	2 mm	Urban_soil	2	5.0	4.16	16.8	0.09
D11	2 mm	Urban_soil	2	5.1	3.89	23.7	0.14
D12	2 mm	Urban_soil	2	5.1	4.13	19.0	0.11
D13	0.2 mm	Urban_soil	2	5.0	3.93	21.4	0.12
D14	0.2 mm	Urban_soil	2	5.0	3.97	20.6	0.12
D15	0.2 mm	Urban_soil	2	5.1	4.32	15.3	0.08
D16	0.2 mm	Urban_soil	2	5.1	4.29	15.9	0.09
D17	0.2 mm	Urban_soil	2	5.1	4.16	18.4	0.10
D18	0.2 mm	Urban_soil	2	5.1	4.11	19.4	0.11
E1	0.2 mm	Tropical_rain_forest	2	5.0	4.01	19.8	0.11
E2	0.2 mm	Tropical_rain_forest	2	5.0	3.74	25.2	0.15
E3	0.2 mm	Tropical_rain_forest	2	5.0	3.82	23.6	0.13
E4	0.2 mm	Tropical_rain_forest	2	5.1	3.98	22.0	0.12
E5	0.2 mm	Tropical_rain_forest	2	5.1	3.85	24.5	0.14
E6	0.2 mm	Tropical_rain_forest	2	5.0	3.96	20.8	0.12
E7	2 cm	Tropical_rain_forest	2	5.0	4.00	20.0	0.11
E8	2 cm	Tropical_rain_forest	2	5.0	3.75	25.0	0.14
E9	2 cm	Tropical_rain_forest	2	5.0	4.02	19.6	0.11
E10	2 cm	Tropical_rain_forest	2	5.1	3.62	29.0	0.17
E11	2 cm	Tropical_rain_forest	2	5.1	3.91	23.3	0.13
E12	2 cm	Tropical_rain_forest	2	5.1	3.78	25.9	0.15
E13	2 mm	Tropical_rain_forest	2	4.9	2.95	39.8	0.25
E14	2 mm	Tropical_rain_forest	2	5.16	4.48	13.2	0.07
E15	2 mm	Tropical_rain_forest	2	5.0	4.07	18.6	0.10
E16	2 mm	Tropical_rain_forest	2	5.02	3.94	21.5	0.12

	Treatment	Locality	Time	Before	After	Percentage	k
E17	2 mm	Tropical_rain_forest	2	5.0	3.88	22.4	0.13
E18	2 mm	Tropical_rain_forest	2	5.1	3.87	24.1	0.14
F1	2 cm	Permaculture	2	5.1	3.51	31.2	0.19
F2	2 cm	Permaculture	2	5.1	4.01	21.4	0.12
F3	2 cm	Permaculture	2	5.0	3.88	22.4	0.13
F4	2 cm	Permaculture	2	5.0	4.12	17.6	0.10
F5	2 cm	Permaculture	2	5.1	3.41	33.1	0.20
F6	2 cm	Permaculture	2	5.1	3.13	38.6	0.24
F7	0.2 mm	Permaculture	2	5.1	3.73	26.9	0.16
F8	0.2 mm	Permaculture	2	5.0	3.88	22.4	0.13
F9	0.2 mm	Permaculture	2	5.1	3.97	22.2	0.13
F10	0.2 mm	Permaculture	2	5.1	3.90	23.5	0.13
F11	0.2 mm	Permaculture	2	5.1	3.87	24.1	0.14
F12	0.2 mm	Permaculture	2	5.0	3.50	30.0	0.18
F13	2 mm	Permaculture	2	5.0	4.16	16.8	0.09
F14	2 mm	Permaculture	2	5.1	4.19	17.8	0.10
F15	2 mm	Permaculture	2	5.0	4.15	17.0	0.09
F16	2 mm	Permaculture	2	5.0	4.22	15.6	0.08
F17	2 mm	Permaculture	2	5.0	4.22	15.6	0.08
F18	2 mm	Permaculture	2	5.0	3.98	20.4	0.11
G1	0.2 mm	Urban_soil	3	5.1	4.13	19.0	0.07
G2	0.2 mm	Urban_soil	3	5.1	4.11	19.4	0.07
G3	0.2 mm	Urban_soil	3	5.0	4.00	20.0	0.07
G4	0.2 mm	Urban_soil	3	5.0	4.36	12.8	0.05
G5	0.2 mm	Urban_soil	3	5.0	3.82	23.6	0.09
G6	0.2 mm	Urban_soil	3	5.0	4.25	15.0	0.05
G7	2 mm	Urban_soil	3	5.1	3.88	23.9	0.09
G8	2 mm	Urban_soil	3	5.0	3.88	22.4	0.08
G9	2 mm	Urban_soil	3	5.0	3.75	25.0	0.10
G10	2 mm	Urban_soil	3	5.0	3.83	23.4	0.09
G11	2 mm	Urban_soil	3	5.0	3.69	26.2	0.10
G12	2 mm	Urban_soil	3	5.0	3.12	37.6	0.16
G13	2 cm	Urban_soil	3	5.0	3.68	26.4	0.10
G14	2 cm	Urban_soil	3	5.1	3.69	27.6	0.11
G15	2 cm	Urban_soil	3	5.0	3.88	22.4	0.08
G16	2 cm	Urban_soil	3	5.1	4.07	20.2	0.08
G17	2 cm	Urban_soil	3	5.0	3.77	24.6	0.09
G18	2 cm	Urban_soil	3	5.1	3.78	25.9	0.10
H1	2 cm	Tropical_rain_forest	3	5.0	3.47	30.6	0.12
H2	2 cm	Tropical_rain_forest	3	5.0	3.18	36.4	0.15
H3	2 cm	Tropical_rain_forest	3	5.1	3.10	39.2	0.17
H4	2 cm	Tropical_rain_forest	3	5.0	3.19	36.2	0.15
H5	2 cm	Tropical_rain_forest	3	5.1	3.32	34.9	0.14
H6	2 cm	Tropical_rain_forest	3	5.1	3.80	25.5	0.10
H7	2 mm	Tropical_rain_forest	3	5.1	3.30	35.3	0.15

	Treatment	Locality	Time	Before	After	Percentage	k
H8	2 mm	Tropical_rain_forest	3	5.0	3.60	28.0	0.11
H9	2 mm	Tropical_rain_forest	3	5.1	3.74	26.7	0.10
H10	2 mm	Tropical_rain_forest	3	5.1	3.81	25.3	0.10
H11	2 mm	Tropical_rain_forest	3	5.1	3.61	29.2	0.12
H12	2 mm	Tropical_rain_forest	3	5.0	3.44	31.2	0.12
H13	0.2 mm	Tropical_rain_forest	3	5.0	3.56	28.8	0.11
H14	0.2 mm	Tropical_rain_forest	3	5.1	3.51	31.2	0.12
H15	0.2 mm	Tropical_rain_forest	3	5.1	3.59	29.6	0.12
H16	0.2 mm	Tropical_rain_forest	3	5.1	3.53	30.8	0.12
H17	0.2 mm	Tropical_rain_forest	3	5.1	3.44	32.5	0.13
H18	0.2 mm	Tropical_rain_forest	3	5.1	3.44	32.5	0.13
I1	2 cm	Permaculture	3	5.0	3.53	29.4	0.12
I2	2 cm	Permaculture	3	5.0	3.54	29.2	0.12
I3	2 cm	Permaculture	3	5.0	3.38	32.4	0.13
I4	2 cm	Permaculture	3	5.0	3.37	32.6	0.13
I5	2 cm	Permaculture	3	5.1	3.66	28.2	0.11
I6	2 cm	Permaculture	3	5.0	3.31	33.8	0.14
I7	0.2 mm	Permaculture	3	5.0	3.83	23.4	0.09
I8	0.2 mm	Permaculture	3	5.0	3.72	25.6	0.10
I9	0.2 mm	Permaculture	3	5.1	3.58	29.8	0.12
I10	0.2 mm	Permaculture	3	5.0	3.65	27.0	0.10
I11	0.2 mm	Permaculture	3	5.1	3.48	31.8	0.13
I12	0.2 mm	Permaculture	3	5.0	3.40	32.0	0.13
I13	2 mm	Permaculture	3	5.1	3.80	25.5	0.10
I14	2 mm	Permaculture	3	5.1	3.57	30.0	0.12
I15	2 mm	Permaculture	3	5.1	3.66	28.2	0.11
I16	2 mm	Permaculture	3	5.1	3.55	30.4	0.12
I17	2 mm	Permaculture	3	5.1	3.55	30.4	0.12
I18	2 mm	Permaculture	3	5.1	3.55	30.4	0.12